

Policy brief on potential targets to reduce risks for health and ecosystems

Draft document prepared by the Task Force on Integrated Assessment Modelling and the Centre for Integrated Assessment Modelling, 9 November 2023

Summary

At its 61st meeting the Working Group on Strategies and Review took note of the information presented by the Task Force on Integrated Assessment Modelling the EMEP Centre for Integrated Assessment Modelling on the feasibility of introducing a risk-based overarching goal for the Convention, in particular a health damage reduction target. The working group requested to provide a policy brief on the potential implications of introducing collective risk-based targets for the UNECE region to address air pollution impacts on health and ecosystems (work plan item 2.1.12). The present informal document provides a draft version of this policy brief. The document focusses on the attainability of an illustrative 50% reduction target of health risks due to exposure to particulate matter and ozone. The parties are invited to take note of the results and send comments to the secretariat before February 3rd, 2024 in order to enable the Task Force to submit a new version for the 62nd meeting of the Working Group on Strategies.

I. Introduction

1. This report describes policy scenarios up to 2050, as calculated with GAINS for the UNECE region, including EECCA-countries (including Kazakhstan, Kyrgyzstan, Uzbekistan, Turkmenistan, Tajikistan) and West-Balkan countries. The scenarios cover options to address particulate matter and ozone precursors, including methane and the potential policy targets that would be attainable.
2. Improvements in the GAINS and EMEP model include local scale modelling, health impacts assessment methods, as well as the inclusion of soil NO_x emissions and a consistent representation of the condensable fraction of PM. GAINS has been prepared to assess sectoral policies (and ‘staged approaches’). The cost-optimized scenarios cover the whole UNECE region.

II. Policy scenarios

3. Three scenarios were developed by the Centre for Integrated Assessment Modelling (CIAM):
 - a. A baseline scenario, considering trends and policies included in established national air pollution control programmes and, for the EU, the European Green Deal including the ‘Fit for 55’ legislation package. For countries without such plans, IEA and FAO projections were used. The baseline includes air pollutants (SO₂, NO_x, PM_{2.5}, NH₃, NMVOC, as well as Black Carbon) and methane emissions up to 2050 and assumes effective implementation of current legislation (CLE).
 - b. A maximum technically feasible reductions (MTFR) scenario uses the same activity data as the baseline and includes implementation of technologies with lowest emission factors in the GAINS model database. These control options include measures to reduce ammonia emissions from agriculture, measures to reduce PM_{2.5} and non-methane volatile organic compounds (NMVOC) emissions from residential solid fuel burning and agricultural waste burning, mitigation technologies to reduce emissions of sulfur dioxide (SO₂), nitrogen oxides (NO_x), and PM_{2.5} for industrial combustion, process and transportation sources, measures to reduce NMVOC from solvent use, and liquid fuel’s storage and distribution, as well as measures to reduce methane (CH₄) emissions from municipal waste treatment, the fossil fuel sector and agriculture. Maritime emission control areas or initiatives by port authorities are assumed to encourage clean ships and to provide shore-to-ship electricity access. The MTFR scenario uses


information about the age structure of installations. Early shutdown or scrapping of cars or boilers is not assumed. Consequently, the mitigation potential increases towards 2050.

- c. An alternative 'LOW' scenario, that includes climate policies compatible with the Paris Agreement goals for the whole world, MTRF measures (also for maritime shipping), improvements of nitrogen use efficiency in agriculture, and dietary changes, towards lower meat protein consumption.
4. The baseline scenario shows strong reductions of air pollutants between 2005 and 2030 (SO₂: -80%, NO_x: -50 to -80%, PM_{2.5}: -25 to -70%) in the EU, North America, and also in West Balkan countries, owing to the Energy Community agreements that include commitments to strong reduction of emissions from stationary sources in the coming decades. Fossil fuel use in EECCA countries continues to grow, however, due to ongoing technical progress, emissions of SO₂ and NO_x are expected to be reduced over time, by approximately 40% and 20%, respectively, between 2005 and 2030. For NH₃ current abatement policies are very modest and the estimated reductions, if any, are primarily due to projected decline in livestock numbers in some regions.
 5. The MTRF scenario shows that for SO₂ most of further mitigation potential is committed in the current legislation, except for EECCA. Similar picture for NO_x, although more further potential available. For NH₃, the mitigation potential is similar across all regions, however, compared to the baseline, the overall potential smaller than that for other air pollutants. Large further mitigation exists for PM_{2.5}, except the EU+EFTA region, especially in EECCA and West Balkan. LOW scenario shows that strong climate action brings additional air pollutant reduction although it is most significant for NH₃ and CH₄ (the latter not shown) and it is in fact due to modelled dietary changes and strong improvements of nitrogen use efficiency in agriculture.
 6. Analysis done for West Balkan and EECCA shows that the local contribution of residential combustion is dominating in many cities. The power sector is an important regional source. In West Balkan, local residential heating sources cause 50% or more of the concentrations. Even, in cases where the baseline brings reductions, the future levels of pollution remain well above the WHO guidelines. This points to the need to develop further mitigation strategies that address both local, regional and transboundary sources to achieve significant reductions of the impact of air pollution in cities in the future.
 7. Methane declines in the baseline only in the EU (due to the Green Deal). This contributes to a 7% methane emission reduction in the UNECE region (excluding North America) between 2015 and 2050. In the rest of the world an emission increase of 33% is expected, associated with the growth of the fossil fuel sector. There is a significant technical emission reduction potential, especially with measures in the fossil fuel and waste sector. This could result in a 63% reduction in the UNECE (excluding North America) and of 33% reduction in the rest of the world between 2015 and 2050. Combined with dietary change, the LOW scenario could reach a reduction of 77% in the UNECE and of 40-45% in the rest of the world.

Figure 1: Emission trends in baseline (CLE), MTR and LOW scenario


Sulfur dioxide (SO₂)

Nitrogen oxides (NO_x)



Ammonia (NH₃)


Particulate matter (PM_{2.5})



III. Impacts for health and ecosystems

8. Calculations with the GAINS model show that most of the population in the UNECE domain (excl. North America) lives in areas where $PM_{2.5}$ is above the current WHO annual mean guideline value of $5 \mu\text{g}/\text{m}^3$. The baseline scenario causes declining concentrations in the EU. The current EU limit value ($25 \mu\text{g}/\text{m}^3$) will be met in 2030. Still elevated concentrations persist in Balkan and EECCA countries (see figure 2). Overall levels in large parts of the EMEP domain remain above the WHO guideline in 2030. The MTR scenario for 2030 does not bring a lot of improvement in the number of people exposed to exceedances of the WHO guideline, although the concentrations and associated health impacts drop. Both MTR and LOW are not yet fully effective in 2030 due to the short time available for full introduction of abatement measures or transformations embedded in the LOW scenario.

Figure 2: Population exposure to $PM_{2.5}$ in the UNECE (excluding North America)




9. The baseline for 2050 shows further improvements, yet the WHO guideline level would only be attained for 1/3 of the population. MTR brings large scale improvements, also across the Balkan, as there is enough time to introduce further technical measures. Finally, the LOW scenario gives even lower concentrations. More than 60% of the population in the UNECE (excluding North America) would be exposed to $PM_{2.5}$ levels above the WHO guideline by 2050 (over 80% in the EU+EFTA+UK, 30% in EECCA + Türkiye). Nearly 30% is exposed to more than $10 \mu\text{g}/\text{m}^3$.

10. For the EU, the exceedance of the critical loads for acidification will be reduced in the baseline scenario from about 9% of all ecosystems in 2015 to 3% in 2030 and 2% in 2050. In the LOW scenario, the exceedance in the EU could drop to below 1% of the ecosystems by 2050. For non-EU countries in the EMEP domain, the exceedance will decline from about 4% of the ecosystems in 2015 to 2% in the 2050 baseline and less than 0.5% in the LOW scenario.

11. The exceedance of the critical loads for eutrophication in the EU will be reduced in the baseline scenario from 80% of all ecosystems in 2015 to 70% in 2030 and 65% in 2050. In the LOW scenario, the area with exceedances could be more than 50% less than in 2015, and 35% of the ecosystems in the EU will remain with an exceedance, even in 2050. For non-EU countries in the EMEP domain, the exceedance will decline from 50% of the ecosystems in 2015 to around 43% in the 2050 baseline and 15% in the LOW scenario.

Figure 3: Exceedance of critical loads for acidification and eutrophication in Europe



IV. Options for policy targets


12. One of the Saltsjöbaden workshop recommendations is to formulate a common target for the air pollution related health risks. Halving the pollution related mortality was suggested. Could this be feasible for the UNECE-region? Can the target be the same for all parties? Can a target be applied to PM_{2.5} related mortality, as well as to mortality due to other pollutants, such as ozone? Should the target focus on mortality, or also include morbidity?

13. Several factors influence the attainability of a 50% reduction target:

- a) What is the base year and what is the target year? Using 2005 as base year will for many countries make the attainability easier than using more recent years, as improvements after 2005 can be taken into account. However, also the choice of the target year influences the attainability.
- b) Should the target be applied to the UNECE as a whole or to each country (or even each city)? Obviously, it is easier to meet the target for a larger area than for each densely populated area.
- c) An important factor is definition of the indicator: do we want to halve the absolute number of attributable deaths? Or do we want to halve the attributable mortality (i.e. deaths per 100.000 inhabitants)? The absolute number will be influenced by population growth between 2005 and 2050, which makes it harder to meet the 50% target.
- d) Should the target be formulated for PM_{2.5} only, or for the combined effect of air pollutants? Inclusion of ozone would make the attainability harder due to the increasing emissions of ozone precursors, such as methane.
- e) The choice of the health impact assessment method, will also influence the attainability. Do we want to include the risks of natural PM, or focus the target on the avoidable (anthropogenic) PM-exposure? The impact of adding morbidity for the attainability of a 50% reduction target would require further analysis.


14. Figure 4 shows the scope for reducing the average exposure to anthropogenic PM_{2.5} in the UNECE as a whole as estimated with the GAINS model. In the baseline scenario a reduction of 50% compared to the 2005 level can already be met in 2030. A 50% reduction compared to the 2015 level would require additional efforts.

Figure 4: Average exposure to PM2.5 in the UNECE-region (incl. North America)




15. In figure 5 the absolute number of premature deaths attributable to PM2.5 exposure is shown. Due to population increase and aging absolute numbers tend to increase, especially in EECCA countries, Türkiye and North America. This makes it impossible to meet the 50% target with baseline policies. Annex 1 shows the differences in attainability of a 50% reduction for absolute numbers across the UNECE region, that are for a large part due to population dynamics.

Figure 5: Absolute numbers of annual premature deaths due to PM2.5 exposure in the UNECE-region (incl. North America)



16. The health risk indicator (premature deaths per 100.000 inhabitants) shows that a 50% reduction target (from the 2005 level) can almost (but not completely) be reached with existing baseline policies (see figure 6). This risk approach would make the health target easier to attain than a 50% reduction of absolute numbers.

Figure 6: Mortality risk due to PM2.5 (annual premature deaths per 100.000) in the UNECE-region (incl. North America)



17. For some countries, 50% reduction of absolute premature death numbers between 2015 and 2050 would not be feasible, even with the LOW scenario. The GAINS model has been made ready to compute alternative, justifiable approaches, such as the least cost outcome to meet the health target for the UNECE as a whole (excluding North America), or an approach that requires an equal reduction of the gap between baseline and MTFR by parties. The following paragraphs describe possible outcomes of such approaches. Note that they are preliminary and only meant as illustrations of the available modelling tool.


18. Full enforcement of baseline policies will achieve approximately 20% reduction of premature deaths by 2050 compared to 2015 when considering population growth and aging. A UNECE-wide 70% gap closure of the range between baseline and MTFR would be sufficient to meet the 50% reduction target. An equal 70% gap closure per country would be more equitable. However, this will result in 30% higher costs.

19. Inclusion of additional climate and dietary change policies (as in the LOW-scenario) would achieve over half of the emission reduction needed to reach 50% reduction in absolute numbers of premature deaths, compared to the baseline scenario. Additional air pollution control costs would be over ten times lower (see figure 7). In either case, some countries are not achieving the 50% target or even show increase in premature mortality compared to 2015.

Figure 7: Cost curves (least cost options) for reducing the number of premature deaths by 50% between 2015 and 2050

Least-cost reduction of PM health impacts in UNECE (excl. North America) by 2050

- Optimization results for UNECE-wide improvements
- Optimization results for equal improvement in all countries




The analysis considers population growth and aging

Source: GAINS model (CIAM)

20. Figure 8 shows illustrative least-cost results of the gap closure approach for the UNECE as a whole (excluding North America) and of a gap closure per country. As a 50% health target will, in the EU, already be met with current legislation, these scenarios would mainly lead to additional costs in non-EU countries and in some countries exceed 0.5% of GDP.

Figure 8: Country outcomes of least cost scenarios to reduce the number of premature deaths by 50% between 2015 and 2050


Results for the 'Baseline' considering population growth and aging, 2050



Source: GAINS model (CIAM/IIASA)

21. Figure 9 shows the least-cost results of the same gap closure approach as applied in Figure 8, but keeping the population constant at 2015 levels. In this case the additional costs for non-EU countries will generally be lower than 0.05% of GDP.

Figure 9: Country outcomes of least cost scenarios to reduce the mortality risks by 50% between 2015 and 2050




V. Options for ozone policy targets

22. In the baseline scenario, average ozone concentrations in Europe will increase by 2-5% between 2015 and 2050. Peak season concentrations will be reduced by around 5-10%. In both cases, the methane emission increase in the baseline scenario hampers the reductions expected from NO_x/VOC reductions within Europe.

23. CIAM estimates that with dietary change bringing reductions in livestock numbers, as included in the 2050 LOW scenario, methane emissions in the UNECE region can be reduced by almost 70% between 2015 and 2050. Combined with a 50% methane emission reduction in the rest of the World compared to 2015, the 2050 LOW scenario would reduce annual mean ozone concentrations by around 15% and peak season concentrations by around 25%. About 20% of the annual mean ozone reduction is driven by reductions in methane, compared to only 12% for peak season concentrations reductions. For mean ozone concentrations, transcontinental non-methane sources dominate over European sources, whilst for peak season concentrations, European NO_x and VOC sources dominate.

24. The difference in ozone concentrations (both ozone mean and ozone peak season) between the 2050 baseline and the 2050 LOW scenario can be attributed for roughly 1/3 to the reduction in global methane emissions, for 1/3 to the reduction of European non-methane precursor emissions and for 1/3 to the reduction of non-methane precursor emissions outside Europe (see figure 10).


Figure 10: Reductions in the maximum 8-hours daily average concentrations (MDA8) in the ozone season



Source: EMEP model (MSC-W)

25. Also for ozone the health impacts depend on whether population dynamics are taken into account, or not. When population growth and aging are taken into account, the absolute number of premature deaths in the UNECE region as a whole (excluding North America) would in the 2050 LOW scenario be 53% lower than in 2015. However, with the mortality risk-based approach, and the assumption of a static population, a decrease of 62% would be possible in the LOW-scenario. Note that the number of premature deaths due to ozone is about 10 times lower than that of $\text{PM}_{2.5}$. Estimates based on the peak-season ozone exposure are preliminary.

Figure 11: Reduction in annual premature deaths due to ozone exposure in the UNECE-region (excl. North America) with expected population growth (right) and without population growth (left)



Source: EMEP (MSC-W) and GAINS (CIAM) models; Split of impacts from UNECE vs global NOx/VOC reductions preliminary and not yet available for 2015 to 2050CLE case. Preliminary results pending further updates to health impact calculation methodology (HRAPIE2 upcoming).

VI. Conclusion

26. A 50% health target (in terms of premature deaths due to PM) appears achievable in the UNECE region as a whole, for most regions (groups of countries) and for many single countries, but not all. Feasibility depends on details of the calculation, reference year, formulation of potential other targets (e.g., for cities, adding morbidity). For the EU, as a whole, the target is already achieved in the baseline scenario. Some non-EU countries may struggle to achieve such a target for themselves. A risk-based target (roughly) proportional to anthropogenic PM_{2.5} exposure seems more achievable. A 50% target for the whole region would be more cost-effective, however less equitable, than the same target for all countries. This preliminary analysis, shows that pursuing additional climate measures and dietary change policies could reduce additional air pollution control costs ten-fold.

27. A 50% target for the reduction of premature deaths due to ozone between 2015 and 2050 will be more challenging. Current air pollution policies are largely offset by the global increase in methane emissions. Contrary to PM_{2.5}, the feasibility of the ozone target is more dependent on global cooperation to reduce ozone precursors, including methane. However, action on methane would only be part of the solution. NO_x and NMVOC emission reductions would still be very important to reduce ground level ozone within the UNECE region.

28. The 2050 LOW scenario would, in EU and non-EU countries, compared to 2015, lead to more than 50% reduction in the area with an exceedance of critical loads, for acidification as well as for eutrophication.

29. Further analysis will consider, inter alia, alternative target setting, including achievement of other health end-point indicators, and inclusion of urban hot-spots in country-based targets. Further work is also needed on validation and improvement of cost estimates and the assessment of cost of non-technical measures.

Annex 1: Attainability of health improvement goals in selected regions

Figure A.1: Change in PM2.5 concentrations and number of premature deaths (taking into account population change and ageing – dynamic demography) for Baseline and LOW scenarios (1)





Figure A.2: Change in PM2.5 concentrations and number of premature deaths (taking into account population change and ageing – dynamic demography) for Baseline and LOW scenarios (2)



Annex 2: Country tables

Table 1: Emissions in GAINS-LRTAP scenarios

SO2 - kt SO2		Baseline		MTR		LOW	
Region	2005	2030	2050	2030	2050	2030	2050
EU-27	6798	794	578	621	322	621	322
EFTA+UK	818	169	131	149	112	104	72
West Balkan	886	181	123	141	44	107	15
EECCA+Turkiye	6530	3629	3938	2714	887	1980	545
North America	15765	2865	2640	2159	1177	1550	959
Total	30796	7639	7411	5783	2542	4362	1914

NOx - kt NO2 (soil NOx is excluded)

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	9694	2463	1331	2161	882	2161	882
EFTA+UK	1960	545	325	484	238	484	205
West Balkan	269	162	130	128	73	114	44
EECCA+Turkiye	6277	4736	5290	3510	1498	3122	950
North America	19025	7637	6589	5853	3455	5180	1843
Total	37225	15543	13666	12137	6146	11062	3923

NOx - kt NO2 (soil NOx included)

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	10470	3236	2061	2680	1372	2631	1196
EFTA+UK	2072	646	426	554	308	556	254
West Balkan	291	186	157	145	92	128	56
EECCA+Turkiye	6720	5295	5906	3902	1928	3459	1201
North America	19627	8342	7313	6161	3768	5463	2049
Total	39180	17705	15864	13441	7467	12237	4756
Increase due soil	105%	114%	116%	111%	122%	111%	121%

NH3 - kt NH3

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	3813	3403	3267	2939	2327	2756	1460
EFTA+UK	391	352	358	309	267	289	177
West Balkan	106	112	117	96	83	81	54
EECCA+Turkiye	1926	2250	2577	1959	1937	1885	1278
North America	4274	4915	5190	3906	3037	3492	1667
Total	10510	11031	11508	9209	7651	8503	4635

NMVOC - kt NMVOC (agricultural VOC are excluded)

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	7451	3832	3192	3333	2236	3333	2236
EFTA+UK	1411	843	775	718	525	720	534
West Balkan	329	326	258	187	64	178	82
EECCA+Turkiye	5390	4755	5073	3466	1934	3245	1829
North America	12764	10171	9705	9746	5295	9554	3772
Total	27345	19928	19003	17450	10054	17030	8453

PM2.5 - kt PM2.5

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	1781	751	544	586	332	546	321
EFTA+UK	167	81	63	73	48	68	44
West Balkan	186	163	136	89	21	79	17
EECCA+Turkiye	1900	1811	1922	942	324	779	233
North America	1925	1132	1162	727	448	701	356
Total	5959	3938	3826	2418	1173	2173	972

Black Carbon - kt BC

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	337	75	38	54	21	63	25
EFTA+UK	37	10	6	7	4	8	4
West Balkan	18	18	14	9	1	9	2
EECCA+Turkiye	226	176	166	88	28	70	12
North America	326	171	170	160	59	160	39
Total	945	450	394	319	114	311	82

CH4 - kt CH4

Region	2005	2030	2050	2030	2050	2030	2050
EU-27	20063	13093	10859	9486	6734	8781	3950
EFTA+UK	4253	2111	1776	1373	1117	1358	695
West Balkan	670	655	711	556	371	462	189
EECCA+Turkiye	29060	28655	30844	12011	9750	11781	6213
North America	29841	40984	42727	27938	23094	27289	15268
Total	83886	85498	86918	51363	41066	49671	26315

Table 2: Population exposed to PM2.5 levels above 5 µg/m³ (million)

	Baseline			MTFR		LOW		GP compliant	
Country	2015	2030	2050	2030	2050	2030	2050	2030	2050
Austria	8,3	7,0	4,0	3,4	0,5	3,3	0,3	6,2	
Belgium	11,3	11,9	10,4	10,9	3,7	10,5	1,1	11,8	
Bulgaria	7,2	6,1	4,9	4,9	0,1	4,0	0,3	6,1	
Croatia	4,2	3,7	2,8	3,5	0,3	3,2	0,0	3,7	
Cyprus	1,2	1,3	1,4	1,3	1,4	1,3	1,4	1,3	
Czech Rep.	10,6	10,2	4,1	3,5	-	2,8	-	9,9	
Denmark	5,7	4,9	0,0	4,5	-	1,7	-	4,6	
Estonia	0,7	0,0	0,0	-	-	-	-	-	
Finland	2,4	1,1	0,8	0,6	-	0,6	0,6	0,9	
France	63,3	45,5	25,5	31,2	16,4	25,1	12,4	40,4	
Germany	81,7	78,3	35,6	36,9	3,6	33,2	2,8	74,3	
Greece	11,2	10,7	9,6	10,4	8,4	9,9	7,1	10,7	
Hungary	9,8	9,2	7,5	8,7	-	8,2	0,9	9,2	
Ireland	4,0	-	-	-	-	-	-	-	
Italy	59,2	56,3	50,4	55,6	47,7	54,7	31,7	55,9	
Latvia	1,9	0,7	0,2	0,2	-	0,2	-	0,5	
Lithuania	2,9	1,4	0,2	0,0	-	0,0	-	1,3	
Luxembourg	0,6	0,7	0,1	-	-	-	-	0,7	
Malta	0,4	0,4	0,4	0,4	0,4	0,4	0,4	0,4	
Netherlands	16,9	17,6	17,5	17,6	9,7	17,6	0,8	17,6	
Poland	38,3	35,6	13,0	10,9	0,0	10,0	-	35,4	
Portugal	10,0	7,0	5,0	5,5	4,3	3,5	1,1	6,9	
Romania	19,9	17,0	13,1	11,5	1,8	10,0	2,0	16,7	
Slovakia	5,4	5,0	2,1	2,0	-	1,7	-	4,9	
Slovenia	2,1	2,0	1,3	1,7	0,1	1,6	0,2	2,0	
Spain	43,1	34,5	28,7	32,3	26,4	29,9	15,4	33,8	
Sweden	5,1	2,7	1,6	1,8	0,4	0,8	0,9	2,4	
EU-27	427,3	370,6	240,4	259,3	125,4	234,2	79,2	357,5	
Albania	2,9	2,9	2,6	2,9	2,3	2,9	1,7	2,9	
Armenia	2,9	2,9	2,7	2,9	2,7	2,9	2,7	2,9	
Azerbaijan	9,5	10,6	11,0	10,5	10,9	10,5	11,0	10,6	
Belarus	9,5	8,9	7,8	5,1	-	4,6	0,6	8,8	
Bosnia-H	3,5	3,4	2,9	3,3	1,2	3,1	0,8	3,4	
Georgia	3,7	3,5	3,2	3,4	2,3	3,4	2,3	3,5	
Iceland	0,1	0,1	0,1	0,1	0,1	-	-	0,1	
Kazakhstan	17,5	19,6	22,6	19,5	21,4	18,8	19,6	19,6	
Kosovo	1,8	1,6	1,2	1,0	0,1	0,9	0,2	1,6	
Kyrgyzstan	5,5	6,6	7,9	6,8	7,7	6,7	7,0	6,6	
North Macedonia	2,1	2,0	1,8	1,9	0,7	1,7	0,6	2,0	
R Moldova	4,1	3,8	3,3	3,8	-	2,3	-	3,8	
Montenegro	0,6	0,5	0,4	0,0	0,0	0,0	0,0	0,5	
Norway	1,3	0,3	0,3	0,3	-	-	0,7	0,5	
Russia	97,7	92,7	84,1	77,6	38,7	69,5	36,4	92,7	
Serbia	8,9	8,4	7,2	6,0	0,0	5,5	0,8	8,4	
Switzerland	7,9	7,7	7,9	6,0	1,4	6,2	3,9	7,6	
Tajikistan	-	-	-	-	-	-	-	-	
Türkiye	78,1	87,7	94,8	86,4	84,8	85,0	79,8	87,7	
Turkmenistan	-	-	-	-	-	-	-	-	
Ukraine	44,7	40,7	35,9	35,4	11,3	32,7	13,1	40,7	
United Kingdom	64,3	55,5	14,1	51,4	9,2	38,7	6,0	55,0	
Uzbekistan									
Non-EU	366,6	359,5	311,9	324,6	194,8	295,4	187,3	358,9	
Total	793,9	730,1	552,3	583,9	320,2	529,6	266,5	716,4	

Table 3: Years of life lost (million)

Country	Baseline			MTFR		LOW		GP compliant	
	2015	2030	2050	2030	2050	2030	2050	2030	2050
Austria	6,1	3,116	2,275	2,644	1,722		1,575	2,737	
Belgium	10,0	5,046	3,664	4,311	2,698		2,339	4,916	
Bulgaria	7,9	4,301	3,181	3,179	1,734		1,567	4,18	
Croatia	5,5	2,453	1,691	1,766	0,989		0,886	2,377	
Cyprus	0,6	0,468	0,452	0,386	0,298		0,284	0,46	
Czech Rep.	10,6	4,756	3,325	3,984	2,306		2,064	4,388	
Denmark	3,0	1,676	1,242	1,416	0,877		0,774	1,66	
Estonia	0,6	0,367	0,297	0,298	0,195		0,171	0,315	
Finland	1,8	1,181	1,055	1,07	0,864		0,825	1,148	
France	41,0	22,266	17,668	19,518	13,726		12,675	21,059	
Germany	56,6	30,446	23,209	26,131	17,233		15,383	28,705	
Greece	11,4	5,753	5,113	4,757	3,708		3,551	5,686	
Hungary	13,6	7,104	4,253	5,317	2,695		2,399	6,933	
Ireland	1,4	0,71	0,582	0,617	0,447		0,399	0,699	
Italy	63,0	31,188	22,830	28,482	17,362		16,024	29,602	
Latvia	1,8	0,949	0,635	0,729	0,397		0,366	0,903	
Lithuania	2,4	1,384	0,982	1,033	0,583		0,522	1,324	
Luxembourg	0,3	0,164	0,119	0,143	0,089		0,081	0,159	
Malta	0,2	0,191	0,199	0,162	0,141		0,137	0,189	
Netherlands	14,2	7,771	5,723	6,721	4,304		3,736	7,529	
Poland	50,2	19,233	13,217	15,877	8,51		7,957	19,147	
Portugal	5,0	3,006	2,707	2,495	1,873		1,859	2,979	
Romania	25,7	12,42	9,290	9,494	5,348		4,793	11,674	
Slovakia	5,9	2,65	1,836	2,09	1,163		1,036	2,571	
Slovenia	2,2	1,232	0,678	0,903	0,431		0,408	1,225	
Spain	26,6	14,696	12,535	14,296	8,839		8,59	14,288	
Sweden	3,0	2,012	1,726	1,818	1,432		1,359	1,972	
EU-27	371	187	140	160	100	0	92	179	
Albania	5	3,493	3,735	2,249	1,118		0,998	3,448	
Armenia	3	2,396	2,406	1,864	1,122		1,116	2,394	
Azerbaijan	6	5,826	6,641	5,183	3,315		2,79	5,823	
Belarus	9	7,519	7,066	5,474	2,668		2,292	7,300	
Bosnia-H	10	6,538	5,724	4,165	1,497		1,17	6,427	
Georgia	3	3,207	3,434	2,572	1,328		1,196	3,203	
Iceland									
Kazakhstan									
Kosovo	4	3,266	2,930	2,077	0,744		0,551	3,206	
Kyrgyzstan									
North Macedonia	3	2,232	2,065	1,473	0,697		0,558	2,198	
R Moldova	4	3,018	2,845	2,128	1,148		0,876	2,878	
Montenegro	1	0,792	0,686	0,514	0,195		0,155	0,778	
Norway	1	0,786	0,691	0,724	0,576		0,686	0,807	
Russia	90	91,9	89,376	64,848	35,76		30,04	91,664	
Serbia	15	13,893	9,672	9,011	3,186		2,509	13,682	
Switzerland	5	3,111	2,620	2,78	2,028		1,892	2,896	
Tajikistan									
Türkiye	107	80,857	63,154	62,187	32,429		28,423	80,346	
Turkmenistan									
Ukraine	43	35,293	37,629	24,838	14,043		11,562	35,025	
United Kingdom	40	20,374	14,758	17,546	11,18		10,044	19,915	
Uzbekistan									
Non-EU	350	285	255	210	113	0	97	282	0
Total	721	471	396	369	213	0	189	461	0

Table 4: Premature deaths from ozone (cases/yr)

Country	Baseline			MTR		LOW		GP compliant	
	2015	2030	2050	2030	2050	2030	2050	2030	2050
Austria	475	319	336	284	275	279	254	316	
Belgium	362	272	299	252	261	248	246	271	
Bulgaria	596	431	344	373	268	360	244	430	
Croatia	312	197	166	172	130	169	120	196	
Cyprus	54	60	86	55	74	55	71	60	
Czech Rep.	537	384	342	338	274	332	254	382	
Denmark	131	110	110	100	93	99	88	109	
Estonia	23	17	15	15	12	15	11	17	
Finland	70	62	65	57	54	56	51	62	
France	2206	1787	2004	1635	1752	1614	1654	1779	
Germany	3752	2786	2675	2532	2273	2496	2126	2769	
Greece	669	579	633	525	531	515	502	578	
Hungary	678	474	392	410	306	399	279	472	
Ireland	60	65	89	63	84	63	83	65	
Italy	4695	3522	3630	3251	3106	3239	3001	3506	
Latvia	53	36	28	31	21	31	19	36	
Lithuania	92	66	55	57	41	56	37	66	
Luxembourg	18	13	16	12	13	12	13	13	
Malta	22	23	27	21	25	21	24	23	
Netherlands	382	334	381	307	329	303	308	332	
Poland	1463	1075	1008	933	775	916	715	1071	
Portugal	435	378	421	358	388	357	382	376	
Romania	1247	945	834	812	630	786	575	943	
Slovakia	274	204	196	177	151	173	139	204	
Slovenia	112	79	80	70	64	69	60	79	
Spain	2113	1893	2273	1772	2061	1761	2012	1890	
Sweden	168	132	137	120	116	119	109	132	
EU-27	21001	16244	16644	14734	14106	14542	13375	16176	
Albania	126	129	155	112	120	110	112	128	
Armenia	203	206	246	193	199	191	190	206	
Azerbaijan	245	327	513	295	359	287	336	327	
Belarus	271	202	184	173	135	169	124	201	
Bosnia-H	217	160	152	139	116	134	107	160	
Georgia	206	169	176	157	123	154	115	169	
Iceland	4	5	7	5	7	5	7	5	
Kazakhstan	0	0	0	0	0	0	0		
Kosovo	51	39	34	34	27	33	24	39	
Kyrgyzstan	0	0	0	0	0	0	0		
North Macedonia	140	117	123	104	100	101	91	117	
R Moldova	208	177	188	153	137	147	125	177	
Montenegro	37	30	31	27	25	25	23	30	
Norway	81	74	93	70	85	70	83	74	
Russia	3099	2875	2867	2588	2249	2528	2133	2872	
Serbia	559	406	341	347	258	332	229	405	
Switzerland	387	315	372	289	317	286	296	306	
Tajikistan	0	0	0	0	0	0	0		
Türkiye	2444	2803	4049	2639	3611	2616	3521	2801	
Turkmenistan	0	0	0	0	0	0	0		
Ukraine	2580	1986	1797	1745	1373	1693	1275	1983	
United Kingdom	1238	1168	1360	1102	1234	1094	1192	1161	
Uzbekistan	0	0	0	0	0	0	0		
Non-EU	12098	11188	12689	10172	10474	9975	9983	11160	
Total	33099	27432	29332	24905	24581	24517	23358	27336	

Table 5: Acidification (% of ecosystem area exceeding critical loads)

Country	Ecosystem area [km ²]	Baseline			MTR		LOW		GP compliant	
		2015	2030	2050	2030	2050	2030	2050	2030	2050
Austria	38.901	0,6	0,0	0,0					0,0	
Belgium	15.482	42,1	32,4	28,9	30,1	22,9	27,8	15,4	32,4	
Bulgaria	54.242	0,0								
Croatia	36.341	3,3	0,5	0,3	0,1		0,1		0,5	
Cyprus	1.692									
Czech Rep.	23.831	77,8	19,4	5,4	5,7	0,2	5,4	0,0	16,1	
Denmark	6.657	13,5	1,6	0,6	0,6		0,3		1,2	
Estonia	30.583									
Finland	281	0,4	0,3	0,3	0,3	0,1	0,3	0,0	0,3	
France	176.852	5,5	2,1	1,2	0,8	0,0	0,2		2,1	
Germany	103.401	48,3	21,9	16,4	16,0	6,4	14,2	3,5	21,0	
Greece	77.626	0,6	0,1	0,1	0,1		0,1		0,1	
Hungary	29.969	5,8	2,6	1,8	1,5	0,0	0,3		2,4	
Ireland	16.195	1,1	0,6	0,4	0,3	0,1	0,2		0,4	
Italy	100.954	0,5	0,1	0,1	0,1	0,0	0,1	0,0	0,1	
Latvia	44.142	2,2	0,1		0,1				0,1	
Lithuania	26.331	24,0	14,3	8,6	4,8	0,4	3,2		11,9	
Luxembourg	1.376	13,6	1,3	0,7	0,4		0,4		1,1	
Malta	35									
Netherlands	2.755	72,4	71,2	70,5	70,8	68,9	70,4	56,9	71,2	
Poland	95.931	42,0	5,9	2,1	2,3	0,1	2,1	0,1	5,3	
Portugal	41.903	1,6	0,3	0,3	0,2	0,0	0,3	0,0	0,3	
Romania	109.259	0,8	0,2	0,0	0,0	0,0	0,0		0,1	
Slovakia	26.757	5,3	0,7	0,2	0,2		0,2		0,5	
Slovenia	14.052	0,0	0,0	0,0					0,0	
Spain	251.625	0,6	0,1	0,1	0,1	0,0	0,0		0,1	
Sweden	391.665	3,3	1,2	1,0	1,0	0,7	1,0	0,7	1,2	
EU-27	1.718.839	9,1	3,2	2,1	2,0	0,9	1,7	0,6	3,0	
Albania	19.947									
Armenia										
Azerbaijan										
Belarus	66.499	5,6	1,0	0,8	0,3	0,0	0,2		0,8	
Bosnia-H	36.959	11,1	1,1	0,6	0,1				1,1	
Georgia										
Iceland										
Kazakhstan										
Kosovo	4.693	7,0								
Kyrgyzstan										
North Macedonia	16.846	1,0								
R Moldova	3.773									
Montenegro	9.041									
Norway	320.380	9,4	4,9	3,6	3,8	1,9	3,6	1,3	4,6	
Russia	643.092	0,5	0,3	0,4	0,1	0,0	0,1	0,0	0,3	
Serbia	33.005	17,2	0,4	0,3	0,2	0,0	0,0		0,4	
Switzerland	9.733	16,4	11,4	10,6	10,5	7,7	9,3	3,1	11,0	
Tajikistan										
Türkiye										
Turkmenistan										
Ukraine	97.758	1,2	0,0	0,0	0,0		0,0		0,0	
United Kingdom	75.806	10,6	3,7	2,4	2,2	0,8	1,8	0,1	3,4	
Uzbekistan										
Non-EU	1.337.532	4,4	1,7	1,3	1,2	0,6	1,1	0,3	1,6	
Total	3.056.371	7,0	2,5	1,8	1,6	0,7	1,4	0,5	2,4	

Table 6: Eutrophication (% of ecosystem area exceeding critical loads)

Country	Ecosystem area [km2]	Baseline			MTR		LOW		GP compliant	
		2015	2030	2050	2030	2050	2030	2050	2030	2050
Austria	50.489	65,1	42,6	35,4	28,4	12,5	24,9	1,3	37,4	
Belgium	15.552	65,7	53,0	45,4	50,1	41,1	45,0	29,6	52,0	
Bulgaria	54.322	85,6	72,3	67,1	57,3	39,5	51,6	33,6	71,2	
Croatia	36.411	90,5	79,5	77,1	73,7	61,5	73,5	46,4	78,7	
Cyprus	1.691	100,0	100,0	100,0	100,0	100,0	100,0	100,0	100,0	
Czech Rep.	23.831	96,8	81,5	71,0	69,7	35,7	68,6	5,1	80,1	
Denmark	6.665	100,0	99,7	99,2	99,5	96,7	99,2	63,1	99,6	
Estonia	30.592	46,4	33,4	30,1	29,5	17,7	28,7	10,7	31,3	
Finland	41.047	10,6	2,5	0,9	1,4	0,0	1,2		2,2	
France	176.937	83,5	66,4	61,2	57,3	41,8	51,3	11,2	65,9	
Germany	103.988	79,8	68,0	62,3	61,9	46,3	59,0	27,1	67,2	
Greece	77.844	100,0	99,9	99,9	99,9	99,7	99,9	98,8	99,9	
Hungary	30.007	91,2	73,8	70,1	68,8	64,0	67,9	48,6	71,5	
Ireland	16.776	48,2	44,8	42,8	40,4	32,8	33,6	4,3	42,3	
Italy	105.815	71,5	50,0	45,6	42,5	30,0	41,9	17,8	48,8	
Latvia	44.159	91,4	72,6	60,3	58,8	42,4	53,8	38,2	70,2	
Lithuania	26.352	99,0	97,3	94,2	93,1	68,9	89,9	38,4	97,0	
Luxembourg	1.377	100,0	100,0	98,8	97,2	90,1	96,4	54,4	100,0	
Malta	35	100,0	100,0	100,0	100,0	100,0	100,0	100,0	100,0	
Netherlands	2.976	87,8	77,5	72,2	75,5	57,7	66,2	20,7	77,5	
Poland	95.929	75,8	59,6	50,2	46,8	20,4	47,0	6,8	57,7	
Portugal	42.008	84,2	70,5	69,2	64,3	58,8	64,7	56,7	70,1	
Romania	109.333	93,9	90,4	86,4	83,2	67,6	79,3	43,6	90,1	
Slovakia	26.799	96,2	86,9	84,0	82,7	59,5	79,9	32,2	86,4	
Slovenia	14.066	86,1	62,1	57,0	54,9	38,2	52,4	25,2	60,5	
Spain	251.922	95,2	90,4	89,1	85,4	78,3	84,7	72,9	90,0	
Sweden	58.643	14,3	12,5	11,3	11,5	4,8	11,2	2,4	12,5	
EU-27	1.445.569	80,2	69,2	65,0	62,2	49,4	60,0	35,5	68,2	
Albania	19.971	92,9	89,5	89,5	85,1	77,7	85,7	74,9	89,4	
Armenia										
Azerbaijan										
Belarus	66.500	100,0	99,8	99,8	98,4	89,1	96,6	46,0	99,8	
Bosnia-H	37.044	74,5	70,5	69,3	66,6	57,6	65,6	50,4	70,2	
Georgia										
Iceland										
Kazakhstan										
Kosovo	4.703	83,9	69,0	66,5	51,7	38,5	39,5	18,5	67,8	
Kyrgyzstan										
North Macedonia	16.892	83,0	71,2	69,1	63,0	55,1	58,2	51,2	71,1	
R Moldova	3.774	99,8	98,4	98,4	80,6	65,0	76,8	52,5	98,4	
Montenegro	9.059	60,6	52,9	48,8	43,5	36,1	36,7	32,1	52,5	
Norway	303.446	11,9	7,2	5,4	5,4	2,0	4,9	0,4	6,9	
Russia	643.119	50,4	44,1	42,2	33,0	12,1	28,7	5,6	43,9	
Serbia	33.064	91,4	86,1	84,0	78,3	64,1	67,2	42,1	85,8	
Switzerland	24.248	57,6	48,4	45,0	44,4	35,1	40,3	13,0	46,2	
Tajikistan										
Türkiye										
Turkmenistan										
Ukraine	97.773	100,0	100,0	99,9	99,4	96,0	98,8	73,1	99,9	
United Kingdom	71.070	25,6	15,4	11,4	11,1	3,9	9,0	0,4	14,7	
Uzbekistan										
Non-EU	1.330.663	49,6	44,2	42,5	37,4	24,3	34,4	15,4	44,0	
Total	2.776.232	65,6	57,2	54,2	50,3	37,4	47,7	25,9	56,6	